

- Were any of these tours traumatic tours (i.e. did you experience any mental health problems or symptoms of PTSD from any of these tours)?

Yes No

- Prior to taking the Lariam, did you have any pre-existing mental health or PTSD problems?
- If so:
 - (i) Had you reported these problems?
 - (ii) Had you sought medical help?

On the tours that you were prescribed Lariam:

- Which tour(s) were you prescribed Lariam for?
- Where were you stationed at the time when you received your Lariam prescription?
- Did you have an individual medical assessment before you were prescribed the drug?
- Were you given a group assessment where your anti-malarial was discussed?
- What did your assessment(s) entail?:
- Were you informed of possible alternatives to mefloquine (Lariam)?
- Were you advised of any reason why you might not be suitable for Lariam? If so, please list below:

- Were you informed of any risks or side-effects associated with mefloquine (Lariam)? If so please list below:
- Whether or not you were informed of risks or side-effects at your assessment, were you already aware of them?
- Were you warned that (or did you already know that), should side-effects be experienced, you should stop taking the mefloquine (Lariam) and seek medical advice before the next dose?
- How frequently did you take mefloquine (Lariam)?
- When did you start taking the mefloquine (Lariam)?
- How long did you take mefloquine (Lariam) for?
- What side effects did you experience whilst taking Lariam?
- How soon after taking the first dose of Lariam did you notice these side-effects?
- Did you report these side effects to anyone, if so whom?
- Did the effects of Lariam cease once you stopped taking it?
- If not, what effects persisted?
- Have you continued to suffer from them to date?

